



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Letting $\tau \equiv \tan \theta$ and $\gamma \equiv \cot \varphi$, (3) and (4) may be transformed to

$$(gr^2 - 2hv^2)\tau^2 + 2rv^2\tau + gr^2 = 0,$$

$$gr^2\gamma^2 - 2rv^2\gamma - (2hv^2 - gr^2) = 0.$$

Hence, under the conditions of the problem

$$\tau = \frac{r(v^2 + R)}{2hv^2 - gr^2}, \quad \gamma = \frac{v^2 + R}{gr},$$

where

$$R \equiv +\sqrt{v^4 + 2ghv^2 - g^2r^2}.$$

Consequently

$$\cos \theta = \frac{2hv^2 - gr^2}{v\sqrt{2(2h^2v^2 - ghr^2 + r^2v^2 + r^2R)}}, \quad (5)$$

$$\cos \varphi = \frac{v^2 + R}{v\sqrt{2(gh + v^2 + R)}}, \quad (6)$$

By considering solid angles, it may be readily seen that the mass which is deposited on the circular area of radius r , corresponding to the cone of half-angle θ , is given by $\frac{1}{2}m(1 - \cos \theta) \equiv m_\theta$. Similarly the mass issuing from the upper or φ cone and falling on the same area equals $\frac{1}{2}m(1 - \cos \varphi) \equiv m_\phi$. The total mass on the area πr^2 is, therefore, $m_\theta + m_\phi \equiv m_r$. The total mass distributed over the ring $2\pi r dr$ is $(dm_r/dr)dr$, so that the surface density is given by

$$\sigma_r = \frac{1}{2\pi r} \left(\frac{dm_\theta}{dr} + \frac{dm_\phi}{dr} \right). \quad (7)$$

Assuming $v^4 + 2ghv^2 - g^2r^2 \neq 0$ the derivatives involved in (7) may be formed by the aid of (5) and (6), so that, in general

$$\sigma_r = \frac{m}{4\pi v R} \left\{ \frac{(gr^2 + 2hR)(v^2 + R)^2}{[2(2h^2v^2 - ghr^2 + r^2v^2 + r^2R)]^{\frac{3}{2}}} + \frac{g^2(2gh + v^2 + R)}{[2(gh + v^2 + R)]^{\frac{3}{2}}} \right\}.$$

Special cases:

(a) When $R = 0$, that is, when r attains its superior limit $v/g\sqrt{v^2 + 2gh}$ the θ and φ trajectories coincide, the angles θ and φ become supplementary, and $\varphi = \tan^{-1}(1/v\sqrt{v^2 + 2gh})$.

(b) Taking $g = 32$ ft./sec.², $h = 5,000$ ft., $v = 1,024$ ft./sec., the maximum value of r equals $4,608\sqrt{66} \div 37,435.57$ ft. For $r = 0$, $\sigma = 3,669 \times 10^{-12}$ m. When $2hv^2 - gr^2 = 0$ (*i. e.*, $\theta = \pi/2$) $\sigma = 1,612 \times 10^{-13}$ m. When $R = 6,604$, $\sigma = 1,823 \times 10^{-11}$ m. In the last instance $r \div 37,435.0001$ ft. If the entire mass were distributed uniformly over the maximum circle the surface density would be approximately $2,271 \times 10^{-13}$ m.

MECHANICS.

296. Proposed by C. N. SCHMALL, New York City.


A force F is exerted in moving a horizontal cylinder up an inclined plane by means of a crowbar of length l . If R be the radius of the cylinder, W its weight, φ the inclination of the plane to the horizontal and ψ the inclination of the crowbar to the horizon, show that

$$F = \frac{WR \sin \varphi}{l[1 + \cos(\varphi + \psi)]}.$$

SOLUTION BY A. M. HARDING, University of Arkansas.

The cylinder is in equilibrium under the three forces P , Q , and W . Hence, by Lami's Theorem, $P : Q : W = \sin \varphi : \sin \psi : \sin (\varphi + \psi)$.

$$\therefore P = \frac{W \sin \varphi}{\sin (\varphi + \psi)}, \quad Q = \frac{W \sin \psi}{\sin (\varphi + \psi)}.$$



From the triangle AOB we obtain

$$AB = R \tan \frac{\varphi + \psi}{2} = R \frac{\sin (\varphi + \psi)}{1 + \cos (\varphi + \psi)}.$$

Taking moments about B we obtain

$$F \times l = P \times AB = \frac{WR \sin \varphi}{1 + \cos (\varphi + \psi)};$$

whence

$$F = \frac{WR \sin \varphi}{l[1 + \cos (\varphi + \psi)]}.$$

Solved similarly by CLIFFORD N. MILLS.

NUMBER THEORY.

210. Proposed by ELMER SCHUYLER, Brooklyn, N. Y.

If a and b are relatively prime and $(a + b)$ is even, then $(a - b)ab(a + b) \equiv 0, \pmod{24}$.

SOLUTION BY R. M. MATHEWS, Riverside, California.

The statements a and b relatively prime and $a + b$ even imply

$$a = 2m + 1, \quad b = 2n + 1$$

and

$$a + b \equiv 0 \pmod{2},$$

$$a - b \equiv 0 \pmod{2}.$$

If m and n both odd or both even,

$$a - b \equiv 0 \pmod{4}.$$

If m and n be one odd and the other even,

$$(a + b) \equiv 0 \pmod{4}.$$